arXiv:1202.0772v1 [quant-ph] 3 Feb 2012

Keeking on cavity dynamical Casimir effect via three-level atom

A. V. Dodonov and V. V. Dodonov
Instituto de Física, Universidade de Brasília, PO Box 04455, 70910-900, Brasília, Distrito Federal, Brazil

We consider the scenario in which a damped three-level atom (in ladder- or V- configuration) is coupled to a single cavity mode whose vacuum state is amplified by dint of the dynamical Casimir effect. We obtain approximate analytical expressions and exact numerical results for the time-dependent probabilities, demonstrating that the presence of the third level can enrich the system dynamics and its population can serve as a witness of photon generation from vacuum.

PACS numbers: 42.50.Pq, 32.80.-t, 42.50.Ct, 42.50.Hz

I. INTRODUCTION

The physics of three-level quantum systems ("atoms") interacting with quantized modes of electromagnetic field is very rich, and many special cases were studied in numerous papers (see, e.g., [1–12] and references therein). In the majority of studies, the coefficients of the Hamiltonians describing such systems were assumed time-independent. Time-dependent couplings were considered, e.g., in [13], but under the restriction of adiabatic variations. Here we consider the light-matter dynamics when a three-level atom interacts with a single cavity mode, whose frequency is rapidly modulated according to the harmonical law $\omega_t = \omega_0[1 + \varepsilon \sin(\eta t)]$ with a small modulation depth, $|\varepsilon| \ll 1$. We shall use dimensionless variables, setting $\hbar = \omega_0 = 1$. Such a situation can arise, in particular, if the selected mode describes the evolution of electromagnetic field in a cavity with vibrating walls, and one of the most impressive manifestations can be the so called Dynamical Casimir Effect (DCE), i.e., the photon generation from the initial vacuum state induced by the motion of boundaries [14, 15]. The simplest model describing this effect is based on the Hamiltonian [16]

$$H_c = \omega_t n - i\chi_t \left(a^2 - a^{\dagger 2}\right), \; \chi_t = (4\omega_t)^{-1} d\omega_t/dt \,,$$

where a and $a^\dagger$ are the cavity annihilation and creation operators, and $n \equiv a^\dagger a$ is the photon number operator. If the modulation frequency η is close to the parametric resonance frequency, $\eta = 2(1+x)$ with $|x| \ll 1$, then one can expect an exponential growth of the mean number of photons inside the *empty ideal* cavity [17].

However, the situation can be very different if the field mode interacts with a detector while the cavity walls oscillate. For example, it was shown in [18] that no more than *two photons* can be created in the cavity if the field-atom coupling is much stronger than the modulation depth ε. In view of the recent progress in experiments on simulating DCE [19–21], the detailed study of different detection schemes becomes a timely and important task.

Recently, various regimes of the *two-level* detectors were analyzed in [22–24]. On the other hand, *three-level* detectors can be more realistic [25], besides, they have several advantages [26]. Therefore we consider the evolution of the single-mode cavity field interacting with a three-level atom, whose free Hamiltonian is $H_a = \tilde{E}_1\sigma_{11} + \tilde{E}_2\sigma_{22} + \tilde{E}_3\sigma_{33}$ (where $\tilde{E}_i$ is the ith energy level, $|\mathbf{i}\rangle$ is the atomic energy eigenstate and $\sigma_{ij} \equiv |\mathbf{i}\rangle\langle\mathbf{j}|$). We are interested in the cases where the atom-field interaction modifies severely the atomless DCE. Therefore we assume that the $|\mathbf{1}\rangle \leftrightarrow |\mathbf{2}\rangle$ transition is resonant with the unperturbed cavity frequency, $\tilde{E}_2 = \tilde{E}_1 + 1$. Fig. 1 depicts two atomic level structures we consider here: the ladder- (or Ξ)- configuration and the V- configuration, where $\Omega_2 \equiv \tilde{E}_3 - \tilde{E}_2$ and $\Omega_3 \equiv \tilde{E}_3 - \tilde{E}_1$ are the transition frequencies. We also define the detunings between the cavity unperturbed frequency and the other atomic transition frequency as $\Delta_2 \equiv 1 - \Omega_2$ and $\Delta_3 \equiv 1 - \Omega_3$. The set of constants g, g_2 and g_3 quantifies the atom-field dipolar coupling strengths between the energy levels as shown in Fig. 1. We assume these constants to be real and $\mathcal{O}(g_1) \sim \mathcal{O}(g_2) \sim \mathcal{O}(g_3)$ (although this is not the most general case, such a choice describes the main phenomena in the most simple way) and much smaller than unity. Finally, the light-matter interaction Hamiltonian H_I must be included to obtain the total Hamiltonian $H = H_c + H_a + H_I$ with the corresponding Schrödinger equation $d|\Psi\rangle/dt = -iH|\Psi\rangle$. In this paper we focus on the amplification of the vacuum fluctuations, so we consider only the zero-excitation initial state, $|\Psi(0)\rangle = |\mathbf{1}, 0\rangle$.



FIG. 1: (Color online) Diagram of the atomic configurations.

The questions we try to answer are: 1) how the presence of the third level can influence the number of created photons and 2) how large can be occupation probabilities of different levels (this is important from the point of view of the detection of DCE). In the next two sections we study analytically the unitary dynamics of the field and atom for the two configurations in the absence of any damping. The approximate analytical solutions are compared in Sec. IV with exact results obtained by solving numerically the master equation, which takes into account the damping in atomic degrees of freedom. However, we neglect the dissipation in the field mode, assuming that the cavity quality factor is high enough. The last section contains a discussion of the main results.

II. LADDER-CONFIGURATION

For the case depicted in Fig. 1a we assume the Jaynes–Cummings atom-field interaction $H_I = a(g\sigma_{21}+g_2\sigma_{32})+H.c.$, where $H.c.$ stands for the Hermitian conjugate and we neglected the counter-rotating terms. In the weak modulation case $|\varepsilon| \ll 1$ considered here, one can write $\chi_t \simeq 2q\cos(\eta t)$ with $q \equiv \varepsilon(1+x)/4$. Going to the interaction picture via the transformation $|\Psi(t)\rangle = V(t)|\psi(t)\rangle$, with $V(t) = \exp[-it(\eta/2)(n+\sigma_{33}-\sigma_{11})]$, and performing the Rotating Wave Approximation (RWA), one obtains the approximate Hamiltonian governing the time evolution of $|\psi(t)\rangle$

$$\begin{aligned} H_1 &\simeq (ga\sigma_{21}+g_2a\sigma_{32}-iqa^2+H.c.) \\ &\quad +x(\sigma_{11}-n-\sigma_{33})-\Delta_2\sigma_{33}. \end{aligned}$$

A. Strong field-atom coupling

If the coupling between the field and atom is much stronger than the frequency modulation depth, $|\varepsilon| \ll |g|$, we expand the wavefunction in the Fock basis as

$$\begin{aligned} |\psi(t)\rangle &= \sum_{n=0}^{\infty} e^{inxt}\left(e^{-itx}p_{\mathbf{1}n}|\mathbf{1},n\rangle + p_{\mathbf{2}n}|\mathbf{2},n\rangle\right. \\ &\quad \left. +e^{it(x+\Delta_2)}p_{\mathbf{3}n}|\mathbf{3},n\rangle\right) \end{aligned}$$

and obtain the following differential equations for the probability amplitudes:

$$\dot{p}_{\mathbf{1}n} = -i\sqrt{n}g p_{\mathbf{2}(n-1)} + \mathcal{W}_{\mathbf{1}}(n) \tag{1}$$

$$\dot{p}_{\mathbf{2}n} = -ig\sqrt{n+1}p_{\mathbf{1}(n+1)} - ig_2\sqrt{n}e^{i\Delta_2 t}p_{\mathbf{3}(n-1)} + \mathcal{W}_{\mathbf{2}}(n)$$

$$\dot{p}_{\mathbf{3}n} = -ig_2\sqrt{n+1}e^{-i\Delta_2 t}p_{\mathbf{2}(n+1)} + \mathcal{W}_{\mathbf{3}}(n)\ , \tag{2}$$

where for $\mathbf{i} = \mathbf{1},\mathbf{2},\mathbf{3}$

$$\begin{aligned} \mathcal{W}_{\mathbf{i}}(n) &\equiv q\Big[\sqrt{n(n-1)}e^{-2ixt}p_{\mathbf{i}(n-2)} \\ &\quad -\sqrt{(n+1)(n+2)}e^{2ixt}p_{\mathbf{i}(n+2)}\Big]\ . \end{aligned}$$

For the initial state $|\mathbf{1},0\rangle$ only the states $\{|\mathbf{1},2n\rangle, |\mathbf{2},2n-1\rangle, |\mathbf{3},2n-2\rangle\}$ become populated throughout the unitary evolution ($n \geq 1$), so the analytical analysis is simplified.

In the resonant case, $\Delta_2 = 0$, the solution of Eqs. (1)-(2) for $q = 0$ (i.e., in the stationary cavity) is $p_{\mathbf{1}0}(t) =$ const and for $n \geq 2$

$$p_{\mathbf{1}n} = \frac{\sqrt{n}g}{G_n}\left(A_n e^{-iG_n t} - B_n e^{iG_n t} - C_n\right), \tag{3}$$

$$p_{\mathbf{3}(n-2)} = \frac{\sqrt{n-1}g_2}{G_n}\left[A_n e^{-iG_n t} - B_n e^{iG_n t} + \frac{ng^2C_n}{(n-1)g_2^2}\right] \tag{4}$$

$$p_{\mathbf{2}(n-1)} = A_n e^{-iG_n t} + B_n e^{iG_n t}\, , \tag{5}$$

where $G_n \equiv \sqrt{ng^2+(n-1)\,g_2^2}$ and A_n, B_n, C_n are constants dependent on the initial conditions. Now we substitute Eqs. (3)-(5) back into Eqs. (1)-(2), assuming that A_n, B_n, C_n are slowly varying functions of time [22]. If initially (at $t = 0$) all the coefficients $p_{\mathbf{i}n}$ are equal to zero, except for $p_{\mathbf{1}\mathbf{0}} = 1$, then $\dot{p}_{\mathbf{1}\mathbf{0}} = -q\sqrt{2}e^{2ixt}p_{\mathbf{1}\mathbf{2}}$ and for $n \geq 2$

$$\dot{A}_n e^{-iG_n t} - \dot{B}_n e^{iG_n t} - \dot{C}_n = \frac{G_n}{g\sqrt{n}}\mathcal{W}_{\mathbf{1}}(n) \tag{6}$$

$$\dot{A}_n e^{-iG_n t} - \dot{B}_n e^{iG_n t} + \frac{ng^2\dot{C}_n}{(n-1)\,g_2^2} = \frac{G_n\mathcal{W}_{\mathbf{3}}(n-2)}{g_2\sqrt{n-1}}$$

$$\dot{A}_n e^{-iG_n t} + \dot{B}_n e^{iG_n t} = \mathcal{W}_{\mathbf{2}}(n-1)\, . \tag{7}$$

From these equations we see that by adjusting the resonance shift x [contained in the functions $\mathcal{W}_{\mathbf{i}}(n)$], one can reduce the arguments of some exponentials to zero values, whilst other exponentials will oscillate with large arguments. Thus we can perform the RWA and obtain a smaller set of essential resonantly coupled differential equations. For example, simple solutions arise for $2x = \pm G_2$. Then only four probabilities can be significantly different from zero:

$$|p_{10}|^2 \simeq \cos^2(\nu t), \quad |p_{\mathbf{1}\mathbf{2}}|^2 \simeq \left(\frac{g}{G_2}\right)^2\sin^2(\nu t), \tag{8}$$

$$|p_{\mathbf{2}1}|^2 \simeq \frac{1}{2}\sin^2(\nu t), \quad |p_{\mathbf{3}0}|^2 \simeq \frac{1}{2}\left(\frac{g_2}{G_2}\right)^2\sin^2(\nu t), \tag{9}$$

where $\nu \equiv \sqrt{2}qg/G_2$ and $G_2 = \sqrt{2g^2+g_2^2}$. All other probabilities contain extra factors of the order of $(\varepsilon/g)^2$, so they can be neglected in this approximation. We see that at most two photons can be generated with a significant probability and the third level becomes partially populated. (If $g_2 = 0$, then the results coincide with that obtained in [18].)

A similar effect of an indirect interaction between different energy levels was discovered in [27], where the *coupling constants* depended on the time-dependent cavity length $L(t)$ as $g \sim [L(t)]^{-1/2}$, while the cavity frequency was supposed to be constant. One more analog is the effect of "atomic shaking" in cavities with moving boundaries, studied in [28]. Recently, an analog of DCE in three-level systems with time-dependent Rabi frequencies was considered in [29].

Another interesting effect is the appearance of a new resonance for $x = 0$ (it is forbidden for the two-level resonant atom), due to the presence of the coefficient C_2

in Eq. (3). To see this we keep only the slowly oscillating terms, obtaining the set of equations

$$\begin{aligned} G_2\dot{p}_{\mathbf{10}} &= 2qg\, C_2 \\ g\left(\dot{A}_2 e^{-iG_2 t} - \dot{B}_2 e^{iG_2 t} - \dot{C}_2\right) &= qG_2\left(p_{\mathbf{10}} - \sqrt{6}p_{\mathbf{14}}\right) \\ \dot{A}_2 e^{-iG_2 t} + \dot{B}_2 e^{iG_2 t} &= -\sqrt{6}q\left(A_4 e^{-iG_4 t} + B_4 e^{iG_4 t}\right) \\ g_2\left[\dot{A}_2 e^{-iG_2 t} - \dot{B}_2 e^{iG_2 t} + 2(g^2/g_2^2)\dot{C}_2\right] &= -\sqrt{2}qG_2 p_{\mathbf{32}} \end{aligned}$$

and so on. We see that $p_{\mathbf{1}(n\geq 4)}$ and $p_{\mathbf{3}(n\geq 2)}$ are coupled resonantly to C_2, while C_2 is coupled resonantly to $p_{\mathbf{10}}$, so the states $|\mathbf{1}, 2n\rangle$ and $|\mathbf{3}, 2n\rangle$ $(n = 0, 1, 2, \ldots)$ become populated throughout the evolution and many photons can be generated. On the other hand, the coefficients A_2 and B_2 are approximately uncoupled from $p_{\mathbf{10}}$ and from $A_{n>2}, B_{n>2}$ due to the rapidly oscillating imaginary exponentials, so they remain approximately zero for all times. Therefore, the atomic state $|\mathbf{2}\rangle$ is only virtually populated, and making the approximation $p_{\mathbf{2}n} = 0$ for all times, one gets the constraint $p_{\mathbf{1}(n+2)} \simeq -(g_2/g)\sqrt{(n+1)/(n+2)}p_{\mathbf{3}n}$, as we confirmed by numerical simulations. Notice that for $g_2 = 0$ (i.e., without the third level) one has $\dot{C}_2 = 0$ therefore $C_2(t) = 0$ and photons cannot be generated [18, 22].

In the dispersive case, $|\Delta_2| \gg |g_2|$, one can write $p_{\mathbf{30}} \simeq (g_2/\Delta_2)\, e^{-i\Delta_2 t} p_{\mathbf{21}}$ for a stationary cavity $(q = 0)$. After repeating the steps above we find that for $2x = \delta_2/2 \pm J$ (where $\delta_2 \equiv g_2^2/\Delta_2$ is the dispersive shift due to the third level and $J \equiv \sqrt{\delta_2^2/4 + 2g^2}$) the only nonzero probabilities are

$$\begin{aligned} |p_{\mathbf{10}}|^2 &\simeq \cos^2(q\nu_{\mp}t), \quad |p_{\mathbf{12}}|^2 \simeq \frac{1}{2}\nu_{\mp}^2 \sin^2(q\nu_{\mp}t), \\ |p_{\mathbf{21}}|^2 &\simeq \frac{1}{2}\nu_{\pm}^2 \sin^2(q\nu_{\mp}t), \end{aligned} \qquad (10)$$

where $\nu_{\pm} = \sqrt{1 \pm \delta_2/(2J)}$. Thus, the rate of photon generation and the occupation probabilities are influenced by the level $|\mathbf{3}\rangle$, although it remains effectively unpopulated for all times.

B. Weak field-atom coupling

For a weak field-atom coupling, $|\varepsilon| \gg |g|$, many photons can be generated under the resonance condition $x = 0$. To calculate the accompanying atomic dynamics in this case, it is convenient to use the effective Hamiltonian approach instead of solving the differential equations for the probability amplitudes [23]. We rewrite the time-dependent state $|\psi(t)\rangle$ corresponding to the Hamiltonian H_1 as

$$|\psi(t)\rangle = e^{-iH_1 t}|\psi(0)\rangle = U^{\dagger}\exp\left(-iH_{ef}t\right)U|\psi(0)\rangle, \quad (11)$$

where $|\psi(0)\rangle$ is the initial state and we introduced a unitary operator U to define the effective Hamiltonian $H_{ef} \equiv UH_1U^{\dagger}$. In the resonant regime, $\Delta_2 = 0$, we choose the transformation [24]

$$U = e^{iY},\ Y = a^{\dagger}\left(\xi\sigma_{21} + \xi_2\sigma_{32}\right) + a\left(\xi\sigma_{12} + \xi_2\sigma_{23}\right), \quad (12)$$

where $\xi = 2g/\varepsilon \ll 1$ and $\xi_2 = 2g_2/\varepsilon \ll 1$. Then to the second order in ξ

$$H_{ef} = i\theta\left(a^{\dagger 2} - a^2\right) + iq\xi\xi_2\left(\sigma_{13} - \sigma_{31}\right), \quad (13)$$

where $\theta \equiv q[1 + \xi^2(\sigma_{22} - \sigma_{11}) + \xi_2^2(\sigma_{33} - \sigma_{22})]$ is an operator with respect to the atomic basis. This effective Hamiltonian holds approximately for $t \ll 1/g$ (so the product εt can be greater than unity and several photons can be created from vacuum). Moreover, within the idea of RWA we can neglect the last term in (13), as it is nondiagonal in the atomic energy basis. Then the time-dependent wave function reads $|\psi(t)\rangle = U^{\dagger}\hat{\Lambda}_{\theta}U|\psi(0)\rangle$, where $\hat{\Lambda}_v \equiv \exp[vt(a^{\dagger 2} - a^2)]$ is the squeezing operator with nonzero matrix elements [30]

$$\Lambda_v^{(n)} \equiv \langle 2n|\hat{\Lambda}_v|0\rangle = \mathcal{C}_v^{-1/2}(\mathcal{S}_v/\mathcal{C}_v)^n \frac{\sqrt{(2n)!}}{2^n n!},$$

where $|k\rangle$ denotes the kth Fock state and $\mathcal{C}_v = \cosh(2vt)$, $\mathcal{S}_v = \sinh(2vt)$. Using the property $\hat{\Lambda}_v a\hat{\Lambda}_v^{\dagger} = \mathcal{C}_v a - \mathcal{S}_v a^{\dagger}$ one can show that for the initial state $|\psi(0)\rangle = |\mathbf{1}, 0\rangle$ the probability amplitudes read to second order in ξ

$$\langle \mathbf{1}, 2n|\psi(t)\rangle = [1 - \xi^2(n+1)]\Lambda_{\theta_1}^{(n)} + \xi^2(2n+1)\mathcal{C}_{\theta_2}^{-1}\Lambda_{\theta_2}^{(n)},$$

$$\langle \mathbf{2}, 2n+1|\psi(t)\rangle = i\xi\sqrt{2n+1}(\mathcal{C}_{\theta_2}^{-1}\Lambda_{\theta_2}^{(n)} - \Lambda_{\theta_1}^{(n)}),$$

where $\theta_1 = q(1 - \xi^2)$, $\theta_2 = q(1 + \xi^2 - \xi_2^2)$, while $\langle \mathbf{3}, 2n|\psi(t)\rangle \propto \xi\xi_2$, so it is much smaller and we do not write it here explicitly (other probability amplitudes are exactly zero due to the assumed initial state $|\mathbf{1}, 0\rangle$). Therefore, in the resonant regime the third level is not populated within the time scale $\varepsilon t \sim 1$, but the photon statistics is nevertheless slightly modified due to the presence of ξ_2 in θ_2. We checked these expressions by solving numerically the Schrödinger equation and found a good agreement.

III. V-CONFIGURATION

Now we repeat the steps of the previous section for the atom-field interaction Hamiltonian $H_I = a(g\sigma_{21} + g_3\sigma_{31}) + H.c.$, that describes the atomic configuration depicted in Fig. 1b. Using the transformation $|\Psi(t)\rangle = V(t)|\psi(t)\rangle$ with $V(t) = \exp\left[-it\left(\eta/2\right)\left(n + \sigma_{33} + \sigma_{22}\right)\right]$ we obtain

$$\begin{aligned} H_2 \simeq\ & \left(ga\sigma_{21} + g_3a\sigma_{31} - iqa^2 + H.c.\right) \\ & -x(n + \sigma_{22} + \sigma_{33}) - \Delta_3\sigma_{33}. \end{aligned}$$

For the initial state $|\mathbf{1},0\rangle$ only the states $\{|\mathbf{1},2n\rangle, |\mathbf{2},2n-1\rangle, |\mathbf{1},2n-1\rangle\}$ become populated during the unitary evolution ($n\geq 1$). Writing the wavefunction as

$$|\psi\rangle \;=\; \sum_{n=0} e^{ixnt}\left(p_{\mathbf{1}n}|\mathbf{1},n\rangle + e^{itx}p_{\mathbf{2}n}|\mathbf{2},n\rangle \right. \\ \left. + e^{it(\Delta_3+x)}p_{\mathbf{3}n}|\mathbf{3},n\rangle\right),$$

the differential equations for the probability amplitudes read

$$\begin{aligned} \dot{p}_{\mathbf{1}n} &= -ig\sqrt{n}p_{\mathbf{2}(n-1)} - ig_3\sqrt{n}e^{it\Delta_3}p_{\mathbf{3}(n-1)} + \mathcal{W}_{\mathbf{1}}(n) \\ \dot{p}_{\mathbf{2}n} &= -ig\sqrt{n+1}p_{\mathbf{1}(n+1)} + \mathcal{W}_{\mathbf{2}}(n) \\ \dot{p}_{\mathbf{3}n} &= -ig_3\sqrt{n+1}e^{-it\Delta_3}p_{\mathbf{1}(n+1)} + \mathcal{W}_{\mathbf{3}}(n) \end{aligned}$$

and their solution in the resonant regime, $\Delta_3=0$, and for $q=0$ are

$$p_{\mathbf{1}n} = A_n e^{-iG_n t} + B_n e^{iG_n t}, \tag{14}$$

$$p_{\mathbf{2}(n-1)} = \frac{g\sqrt{n}}{G_n}\left(A_n e^{-iG_n t} - B_n e^{iG_n t} + C_n\right),$$

$$p_{\mathbf{3}(n-1)} = \frac{g_3\sqrt{n}}{G_n}\left(A_n e^{-iG_n t} - B_n e^{iG_n t} - \frac{g^2}{g_3^2}C_n\right), \tag{15}$$

where $G_n \equiv \sqrt{n(g^2+g_3^2)}$. Therefore for *weak modulation* ($|\varepsilon| \ll |g|$) the resonances occur only for $2x=\pm G_n$, resulting in the probabilities

$$|p_{\mathbf{1}0}|^2 \simeq \cos^2(qt), \quad |p_{\mathbf{1}2}|^2 \simeq \frac{1}{2}\sin^2(qt), \tag{16}$$

$$|p_{\mathbf{2}1}|^2 \simeq \left(\frac{g}{G_2}\right)^2 \sin^2(qt), \quad |p_{\mathbf{3}1}|^2 \simeq \left(\frac{g_3}{G_2}\right)^2 \sin^2(qt). \tag{17}$$

Moreover, since $p_{\mathbf{12}}$ does not contain the coefficient C_2 in Eq. (14), the $x=0$ resonance does not appear for the V-configuration. In the dispersive regime, $|\Delta_3| \gg |g_3|$, one can write $p_{\mathbf{31}} \simeq (\sqrt{2}g_3/\Delta_3)e^{-it\Delta_3}p_{\mathbf{12}}$, where $\delta_3 \equiv g_3^2/\Delta_3$, and we find that the resonances occur for $2x=\delta_3 \pm J$ with resulting probabilities

$$\begin{aligned} |p_{\mathbf{1}0}|^2 &\simeq \cos^2(q\nu_\pm t), \quad |p_{\mathbf{1}2}|^2 \simeq \frac{1}{2}\nu_\pm^2 \sin^2(q\nu_\pm t) \\ |p_{\mathbf{2}1}| &\simeq \frac{1}{2}\nu_\mp^2 \sin^2(q\nu_\pm t), \end{aligned} \tag{18}$$

where $J \equiv \sqrt{\delta_3^2 + 2g^2}$ and $\nu_\pm = \sqrt{1\pm\delta_3/J}$. Notice that the expressions (18) are slightly different from the corresponding expressions (10) for the ladder-configuration.

For *strong modulation*, $|\varepsilon| \gg |g|$, we perform the transformation

$$U=e^{iY},\; Y=a^\dagger\left(\xi\sigma_{21}+\xi_3\sigma_{31}\right)+a\left(\xi\sigma_{12}+\xi_3\sigma_{13}\right), \tag{19}$$

where $\xi = 2g/\varepsilon \ll 1$ and $\xi_3 = 2g_3/\varepsilon \ll 1$, to obtain the effective Hamiltonian in the resonant regime (valid for $t \ll g^{-1}$)

$$H_{ef} = i[\theta + q\xi\xi_3\left(\sigma_{23}+\sigma_{32}\right)]\left(a^{\dagger 2}-a^2\right), \tag{20}$$

where $\theta \equiv q[1+\xi^2(\sigma_{22}-\sigma_{11})+\xi_3^2(\sigma_{33}-\sigma_{11})]$. Neglecting non-diagonal atomic terms, we obtain for the initial state $|\mathbf{1},0\rangle$ the following nonvanishing time dependent amplitudes:

$$\begin{aligned} \langle\mathbf{1},2n|\psi(t)\rangle &= \left[1-\left(\xi^2+\xi_3^2\right)(n+1)\right]\Lambda_{\theta_1}^{(n)} \\ &\quad + (2n+1)\left[\xi^2\mathcal{C}_{\theta_2}^{-1}\Lambda_{\theta_2}^{(n)} + \xi_3^2\mathcal{C}_{\theta_3}^{-1}\Lambda_{\theta_3}^{(n)}\right], \end{aligned}$$

$$\begin{aligned} \langle\mathbf{2},2n+1|\psi(t)\rangle &= i\xi\sqrt{2n+1}[\mathcal{C}_{\theta_2}^{-1}\Lambda_{\theta_2}^{(n)} - \hat{\Lambda}_{\theta_1}^{(n)}], \\ \langle\mathbf{3},2n+1|\psi(t)\rangle &= i\xi_3\sqrt{2n+1}[\mathcal{C}_{\theta_3}^{-1}\Lambda_{\theta_3}^{(n)} - \Lambda_{\theta_1}^{(n)}], \end{aligned}$$

where

$$\theta_1 = q(1-\xi^2-\xi_3^2),\; \theta_2 = q(1+\xi^2),\; \theta_3 = q(1+\xi_3^2).$$

As expected for the resonant regime, the third level is populated in this case, and the photon statistics is severely modified as compared to the scenario of resonant two-level atom.

IV. A COMPARISON WITH NUMERICAL CALCULATIONS

We verified that our analytical results are in good agreement with exact numerical calculations. Here we demonstrate a few illustrations. To be closer to realistic experimental conditions, we took into account a possibility of damping in the atomic degrees of freedom, using the Lindblad-type Markovian master equation for the total statistical operator ρ of the atom-field system

$$\dot{\rho} = -i\left[H,\rho\right] + \lambda\mathfrak{D}\left[\sigma_{12}\right]\rho + \mathcal{L}\rho, \tag{21}$$

where $\mathcal{L}\rho = \lambda_2\mathfrak{D}[\sigma_{23}]\rho$ for the ladder-configuration and $\mathcal{L}\rho = \lambda_3\mathfrak{D}[\sigma_{13}]\rho$ for V-configuration. The Lindblad kernel is

$$\mathfrak{D}[O]\rho \equiv (2O\rho O^\dagger - O^\dagger O\rho - \rho O^\dagger O)/2,$$

λ_2 is the damping rate for the transition $|\mathbf{3}\rangle \to |\mathbf{2}\rangle$, whereas λ_3 is that for the transition $|\mathbf{3}\rangle \to |\mathbf{1}\rangle$. Although we do not consider specific atoms, we use convenient values of g and λ coefficients within the validity of our approach and close to experimental reality.

In Fig. 2 we show numerical results, obtained by solving numerically the complete master equation, in regimes where analytical solutions are not available, for the following values of dimensionless parameters: $g=3\times10^{-2}$, $g_2=4\times10^{-2}$, $\varepsilon=10^{-3}$, $\lambda=5\times10^{-4}$ and $\lambda_2=\lambda(g_2/g)^2$. Such a choice agrees with a concrete example of the transmon multi-level qubit considered in [31] (assuming that the noise couples to the atom via the dipolar interaction [32]).

Nonvanishing probabilities $p_{\mathbf{i}n}$ ($\mathbf{i}=1,2,3$) for the resonant case ($\Delta_2=0$) with the resonance shift $2x=G_2$ are shown in Fig. 2a, where we see that the third level is populated and at most two photons are created. The dispersive case ($\Delta_2=10g_2$) with resonance shift $2x=\delta_2/2+J$ is considered in Fig. 2b: as expected, the third level is not populated and p_{21} is slightly bigger than p_{12}, in accordance with the predictions given in Eq. (10). In Fig. 2c we analyze the resonance shift $x=0$ for the case of resonant third level in the absence of damping ($\lambda=\Delta_2=0$). This resonance shift is absent for the resonant two-level atom setup, and its dynamical behavior is currently not known analytically due to coupling of many differential equations. Still, from numerical data we see that for $\varepsilon t>1$ photons are generated exponentially, although with much smaller rate than for the empty-cavity case ($\langle n_0\rangle$ denotes the mean photon number in the atomless case), and the third level is significantly populated, while the second level remains unpopulated for all times, as predicted from our qualitative arguments. Thus the resonant ladder-configuration could be used for verifying photon generation from vacuum, since many photons are generated and the third level population can serve as the witness of the DCE.



FIG. 2: (Color online) Atom-field dynamics for the ladder-configuration and different resonance shifts. **a)** $2x=G_2$. **b)** $2x=\delta_2/2+J$. In (a) and (b) for $\varepsilon t=10$ the curves are (from above): $\boldsymbol{p_{10}}$, $\boldsymbol{p_{20}}$ (coincident with $\boldsymbol{p_{11}}$), $\boldsymbol{p_{21}}$, $\boldsymbol{p_{12}}$ and $\boldsymbol{p_{30}}$ [absent in (b)]. **c)** Behavior of average values for $x=0$ in the strong modulation regime. Q is the Mandel factor of the field mode.

In Fig. 3 we illustrate the exact dynamics for the V-configuration. Fig. 3a shows the behavior of probabilities in the resonant regime ($\Delta_3=0$) and weak modulation ($\varepsilon=10^{-3}$) for $2x=G_2$, while in Fig. 3b we consider the dispersive regime ($\Delta_3=-12g_3$) and the resonance shift $2x=\delta_3+J$ [other parameters are: $g=3\times10^{-2}$, $g_3=4\times10^{-2}$, $\lambda=5\times10^{-4}$, $\lambda_3=\lambda(g_3/g)^2$]. In both cases at most two photons are created as predicted analytically in the absence of damping, and in the dispersive case $\boldsymbol{p_{21}}$ lies slightly above $\boldsymbol{p_{12}}$ in accordance with Eq. (18). In Fig 3c we show the photon distribution in the absence of damping (obtained by tracing out the atomic degrees of freedom) for $x=0$ in the strong modulation regime for parameters $\varepsilon=10^{-2}$, $g=5\times10^{-4}$, $g_3=8\times10^{-4}$ and $\Delta_3=-4g_3$ (so the third level is neither in resonant nor in dispersive regime). For comparison we show the photon number distribution in the absence of the third level ($g_3=0$) to emphasize that the photon statistics is substantially modified by the third level. For example, by using post-selection methods based on detecting the atomic state, novel cavity field states could be engineered [23].



FIG. 3: (Color online) Atom-field dynamics for the V-configuration and different resonance shifts. **a)** $2x=G_2$. For $\varepsilon t=10$ the curves are (from above): $\boldsymbol{p_{10}}$, $\boldsymbol{p_{11}}$, $\boldsymbol{p_{12}}$, $\boldsymbol{p_{30}}$, $\boldsymbol{p_{31}}$, $\boldsymbol{p_{20}}$, $\boldsymbol{p_{21}}$; **b)** $2x=\delta_3+J$. For $\varepsilon t=10$ the curves are (from above): $\boldsymbol{p_{10}}$, $\boldsymbol{p_{11}}$ (coincident with $\boldsymbol{p_{20}}$), $\boldsymbol{p_{21}}$, $\boldsymbol{p_{12}}$. **c)** Photon number distribution for $x=0$ and $\varepsilon t=3.5$ in the strong modulation regime and $\Delta_3=-4g_3$ (without damping, $\lambda=\lambda_3=0$), compared to the two-level atom case ($g_3=0$).

V. CONCLUSIONS

Here we obtained simple analytical expressions for the atom-field dynamics when a three-level atom (see Fig. 1) interacts with a single cavity field mode whose vacuum state is being amplified via the dynamical Casimir effect, and confirmed their validity by exact numerical computations. This study is relevant since the actual atoms in cavity QED and artificial atoms in circuit QED are indeed multi-levels. We found that the third level modifies the resonance frequencies as compared to the two-level case, and the dynamical behavior may be drastically different from the cases of atomless cavity or a two-level atom. For instance, the modulation frequency equal to twice the unperturbed cavity frequency does lead to photon generation from vacuum and occupation of the third level in the ladder-configuration, whereas this modulation frequency is forbidden in the case of two-level resonant atom or V-configuration. Therefore the problem studied here might be useful to detect the Casimir photons by measuring the third-level occupancy or creating different entangled states. Hence the presence of the third level is not actually a drawback but an opportunity for observing a reach dynamical behavior.

Acknowledgments

V.V.D. acknowledges the partial support of CNPq (Brazilian agency).

[1] P. L. Knight and P. W. Milonni, Phys. Rep. **66**, 21 (1980).

[2] H.-I. Yoo and J. H. Eberly, Phys. Rep. **118**, 239 (1985).

[3] S. M. Chumakov, V. V. Dodonov, and V. I. Man'ko, in *Classical and Quantum Effects in Electrodynamics*, edited by A. A. Komar, Proc. Lebedev Phys. Inst. Vol. 176 (Nova Science, Commack, New York, 1988), p. 77.

[4] V. Bužek, J. Mod. Opt. **37**, 1033 (1990).

[5] C. C. Gerry and J. H. Eberly, Phys. Rev. A **42**, 6805 (1990).

[6] Fam Le Kien and A. S. Shumovsky, Int. J. Mod. Phys. B **5**, 2287 (1991).

[7] M. Alexanian and S. K. Bose, Phys. Rev. A **52**, 2218 (1995).

[8] Ying Wu and Xiaoxue Yang, Phys. Rev. A **56**, 2443 (1997).

[9] Jing-Bo Xu and Xu-Bo Zou, Phys. Rev. A **60**, 4743 (1999).

[10] A. B. Klimov and L. L. Sanchez-Soto, Phys. Rev. A **61**, 063802 (2000).

[11] M. Fleischhauer and M. D. Lukin, Phys. Rev. A **65**, 022314 (2002).

[12] A. Messina, S. Maniscalco, and A. Napoli, J. Mod. Opt. **50**, 1 (2003).

[13] M. Janowicz, Phys. Rep. **375**, 327 (2003).

[14] V. V. Dodonov, Phys. Scr. **82**, 038105 (2010).

[15] D. A. R. Dalvit, P. A. Maia Neto, and F. D. Mazzitelli, in *Casimir Physics*, edited by D. Dalvit, P. Milonni, D. Roberts, and F. da Rosa, Lecture Notes in Physics Vol. 834 (Springer, Berlin, 2011), p. 419.

[16] C. K. Law, Phys. Rev. A **49**, 433 (1994).

[17] V. V. Dodonov and A. B. Klimov, Phys. Rev. A **53**, 2664 (1996); G. Plunien, R. Schützhold, and G. Soff, Phys. Rev. Lett. **84**, 1882 (2000); M. Crocce, D. A. R. Dalvit, and F. D. Mazzitelli, Phys. Rev. A **64**, 013808 (2001).

[18] V. V. Dodonov, Phys. Lett. A **207**, 126 (1995).

[19] C. M. Wilson, G. Johansson, A. Pourkabirian, M. Simoen, J. R. Johansson, T. Duty, F. Nori, and P. Delsing, Nature (London) **479**, 376 (2011).

[20] P. D. Nation, J. R. Johansson, M. P. Blencowe, and F. Nori, Rev. Mod. Phys. **84**, 1 (2012).

[21] A. Agnesi, C. Braggio, G. Carugno, F. Della Valle, G. Galeazzi, G. Messineo, F. Pirzio, G. Reali, and G. Ruoso, Rev. Sci. Instr. **82**, 115107 (2011).

[22] A. V. Dodonov, R. Lo Nardo, R. Migliore, A. Messina, and V. V. Dodonov, J. Phys. B **44**, 225502 (2011).

[23] A. V. Dodonov and V. V. Dodonov, Phys. Lett. A **375**, 4261 (2011).

[24] A. V. Dodonov and V. V. Dodonov, Phys. Rev. A **85**, 015805 (2012).

[25] B. Peropadre, G. Romero, G. Johansson, C. M. Wilson, E. Solano, and J. J. García-Ripoll, Phys. Rev. A **84**, 063834 (2011).

[26] D. Pinotsi and A. Imamoglu, Phys. Rev. Lett. **100**, 093603 (2008).

[27] T. Taneichi and T. Kobayashi, J. Phys. Soc. Jpn. **67**, 1594 (1998).

[28] A. M. Fedotov, N. B. Narozhny, and Yu. E. Lozovik, Phys. Lett. A **274**, 213 (2000); N. B. Narozhny, A. M. Fedotov, and Yu. E. Lozovik, Phys. Rev. A **64**, 053807 (2001).

[29] I. Carusotto, M. Antezza, F. Bariani, S. De Liberato, and C. Ciuti, Phys. Rev. A **77**, 063621 (2008).

[30] R. R. Puri, *Mathematical Methods of Quantum Optics* (Springer, Berlin, 2001).

[31] J. Koch, T. M. Yu, J. Gambetta, A. A. Houck, D. I. Schuster, J. Majer, A. Blais, M. H. Devoret, S. M. Girvin, and R. J. Schoelkopf, Phys. Rev. A **76**, 042319 (2007).

[32] M. Boissonneault, A. C. Doherty, F. R. Ong, P. Bertet, D. Vion, D. Esteve, and A. Blais, Preprint arXiv:1111.0203.